



Breast C-D-G

Naturally Enhancing Glucuronidation – A Detoxification System

DESCRIPTION

Breast C-D-G, available exclusively from Medical Wellness Associates, is a natural supplement of D-glucaric acid bound to calcium.

FUNCTIONS

Calcium D-glucarate enhances the body's natural defenses against environmental toxins and excess steroid hormones. Through the actions of its metabolite D-glucaro-1,4-lactone, calcium-d-glucarate indirectly favors glucuronidation, an important system that the body relies upon to deactivate and eliminate toxins and steroid hormones.

Glucuronidation is a conjugation process through which potentially carcinogenic environmental toxins, such as polycyclic aromatic hydrocarbons and nitrosamines, and excess steroid hormones are neutralized. During glucuronidation, the enzyme glucuronosyl transferase catalyzes the conjugation of free carcinogens and steroid hormones to glucuronic acid. The resulting glucuronide-bound toxins and hormones can then be safely excreted in bile and urine. The body also deconjugates glucuronide-bound end products. The release of glucuronide-bound substances is beneficial in some circumstances, for example, when the excretion of steroid hormones is undesirable. Yet, the deconjugation process, which works via the enzyme beta-glucuronidase, can result in the re-release of harmful toxins back into the body. The benefits of calcium D-glucarate are realized indirectly through the action of its metabolite D-glucaro-1,4-lactone, which inhibits the deconjugating enzyme beta-glucuronidase found in both blood and tissue. Inhibition of beta-glucuronidase causes a net increase in glucuronidation, and thereby enhances a primary detoxification pathway in the body. Animal studies have documented a decrease in B-glucuronidase activity in breast, liver, colon, skin, lung, and prostate tissue following oral administration of calcium D-glucarate.

Once ingested, calcium D-glucarate is converted to D-glucaric acid in the presence of gastric HCl. The body

normally maintains an equilibrium between D-glucaric acid and its two metabolites, D-glucaro-1,4-lactone and D-glucaro-6,3-lactone.

Calcium D-glucarate provides a slow-release source of D-glucaro-1,4-lactone. When taken directly, D-glucaro-1,4-lactone is metabolized quite quickly and is excreted from the body usually within 1-2 hours of its administration. Calcium D-glucarate, however, is metabolized slowly, remaining in the body for 5 or more hours after ingestion. This results in maintenance of stable levels of D-glucaro-1,4-lactone for long periods of time.

D-glucaric acid is produced endogenously in small amounts. It is also present in some fruits and vegetables, including oranges, apples, grapefruit, broccoli and other cruciferous vegetables.

INDICATIONS

Breast C-D-G may be a useful dietary supplement for those who wish to supplement their diet with D-glucaric acid as a precursor to D-glucaro-1, 4-lactone.

FORMULA (CDG)

Each capsule contains:

Calcium D-Glucarate 500 mg

SUGGESTED USE

One to two capsules daily as a dietary supplement, or as directed by physician.

SIDE EFFECTS

No adverse effects have been reported.

STORAGE

Store in a cool, dry place, away from direct light. Keep out of reach of children.

REFERENCES

Abou-Issa H, Moeschberger M, el-Masry W, et al. Relative efficacy of glucarate on the initiation and promotion phases of rat mammary carcinogenesis. *Anticancer Res* 1995;15(3):805-10.

Abou-Issa H, Koolemans-Beyen A, Meredith TA, Webb TE. Antitumor

(continued on reverse)

synergism between non-toxic dietary combinations of isotretinoin and glucarate. *Eur J Cancer* 1992;28A(4-5):784-8.

Abou-Issa HM, Duruibe VA, Minton JP, et al. Putative metabolites derived from dietary combinations of calcium glucarate and N-(4-hydroxyphenyl)retinamide act synergistically to inhibit the induction of rat mammary tumors by 7,12-dimethylbenz[a]anthracene. *Proc Natl Acad Sci USA* 1988;85(12):4181-4.

Dwivedi C, Heck WJ, Downie AA, et al. Effect of calcium glucarate on beta-glucuronidase activity and glucarate content of certain vegetables and fruits. *Biochem Med Metab Biol* 1990;43(2):83-92.

Dwivedi C, Oredipe OA, Barth RF, et al. Effects of the experimental chemopreventive agent, glucarate, on intestinal carcinogenesis in rats. *Carcinogenesis* 1989;10(8):1539-41.

Heerdt AS, Young CW, Borgen PI. Calcium glucarate as a chemopreventive agent in breast cancer. *Isr J Med Sci* 1995;31:101-105.

Oredipe OA, Barth RF, Dwivedi C, Webb TE. Dietary glucarate-mediated inhibition of diethylnitrosamine-induced hepatocarcinogenesis. *Toxicology* 1992;74(2-3):209-22.

Oredipe OA, Barth RF, Hanausek-Walaszek M, et al. Effects of calcium

glucarate on the promotion of diethylnitrosamine-initiated altered hepatic foci in rats. *Cancer Lett* 1987;38(1-2):95-9.

Oredipe OA, Barth RF, Dwivedi C, Webb TE. Chemopreventive activity of dietary glucarate on azoxymethane-induced altered hepatic foci in rats. *Res Commun Chem Pathol Pharmacol* 1989;65(3):345-59.

Walaszek Z, Szymraj J, Narog M, et al. Metabolism, uptake, and excretion of a D-glucaric acid salt and its potential use in cancer prevention. *Cancer Detect Prev* 1997;21(2):178-90.

Walaszek Z, Hanausek-Walaszek M, Webb TE. Dietary glucarate-mediated reduction of sensitivity of murine strains to chemical carcinogenesis. *Cancer Lett* 1986;33(1):25-32.

Walaszek Z, Hanausek M, Sherman U, Adams AK. Antiproliferative effect of dietary glucarate on the Sprague-Dawley rat mammary gland. *Cancer Lett* 1990;49(1):51-7.

Webb TE, Pham-Nguyen MH, Darby M, Hamme AT 2nd. Pharmacokinetics relevant to the anti-carcinogenic and anti-tumor activities of glucarate and the synergistic combination of glucarate:retinoid in the rat

**These statements have not been evaluated by the Food and Drug Administration.
This product is not intended to diagnose, treat, cure, or prevent any disease.**

**Medical Wellness Associates
6402 Route 30
Jeannette, PA 15644
1-800-834-4325
www.vitamincoach.com**